


CAUTIONARY & ADVISORY LABELS

1 This medicine may cause drowsiness and may increase the effects of alcohol. If affected, do not drive a motor vehicle or operate machinery.	1a This preparation is to aid sleep. Drowsiness may continue the following day. If affected, do not drive or operate machinery. Avoid alcohol.	2 DO NOT TAKE ALCOHOL while being treated with this medicine.	3a Take on an empty stomach at least half an hour before meals and at bedtime.	3b Take on an empty stomach at least half an hour before food or two hours after food.
4a Do not take dairy products, antacids or mineral supplements within two hours of each dose of this medicine.	4b Ask your doctor or pharmacist before taking medicines for heartburn, reflux or indigestion.	5 Ask your doctor or pharmacist before using any other medicine including over-the-counter medicines or health products.	6 REFRIGERATE Do not freeze	7a Discard contents after / /
7b Discard days after opening. Date opened / /	8 Avoid excessive skin exposure to sunlight and sunlamps while being treated with this medicine.	9 DO NOT STOP TAKING THIS MEDICINE ABRUPTLY unless otherwise advised by your doctor.	10a Do not take more than one aspirin tablet or capsule each day while being treated with this medicine.	10b DO NOT TAKE ASPIRIN OR ANTI-INFLAMMATORY MEDICINES while being treated with this medicine unless advised by your doctor.
11 DO NOT TAKE POTASSIUM while being treated with this medicine unless advised by your doctor	12 This medicine may affect mental alertness and/or coordination. If affected, do not drive a motor vehicle or operate machinery.	13 Do not remove from original packaging until dose required.	14 RINSE MOUTH with water after each use.	15a This medicine replaces Do not use both.
15b Active Ingredient: This medicine replaces: DO NOT USE BOTH	16 This medicine may cause dizziness especially when you stand up quickly. Ask your doctor or pharmacist for advice.	18 Avoid eating grapefruit or drinking grapefruit juice while being treated with this medicine.	19a Contains PARACETAMOL. Consult your doctor or pharmacist if taking other paracetamol products.	19b Consult your doctor or pharmacist before taking other medicines for pain or inflammation.
20 Take once weekly on the same day.	21 Special handling and disposal required — ask your pharmacist.	22 Use only with approved or recommended device.	23 This product has been compounded by the pharmacist.	24 FOR 3 DAYS USE ONLY can cause addiction
A SWALLOW WHOLE Do not crush or chew	B TAKE WITH OR SOON AFTER FOOD	C TAKE AT LEAST HALF AN HOUR BEFORE FOOD	E CONTINUE FOR 14 DAYS AFTER SYMPTOMS CEASE	F TAKE IMMEDIATELY BEFORE FOOD
G TAKE IN THE MORNING drink plenty of water	H STORE FROZEN	I CERTAIN FOODS AND JUICES SHOULD BE AVOIDED	J SHAKE WELL BEFORE EACH USE	K FOR EXTERNAL USE ONLY
L CAUTION NOT TO BE TAKEN	M DO NOT SWALLOW	N CONTAINS PEANUT OIL	 <p>Pharmaceutical Society of Australia</p> <p>© Pharmaceutical Society of Australia, reproduced by StirlingFildes with permission.</p> <p>The information in this catalogue chart is no substitute for professional knowledge or judgement and users should exercise their own skill and care with respect to its use. Users are encouraged to consult with other sources, such as the Australian Pharmaceutical Formulary and Handbook and the manufacturer's Product Information to ensure the information provided in this catalogue chart is suitable for their purpose.</p>	

ADDITIONAL LABELS

61 KEEP OUT OF THE REACH OF CHILDREN	62 LAST REPEAT FOR FURTHER REPEATS A NEW PRESCRIPTION IS NECESSARY	62a LAST REPEAT	67 DELIVER	70 The Mixture SHAKE THE BOTTLE
71 Shake the Bottle	98 DO NOT TAKE THIS MEDICINE IF ALLERGIC TO PENICILLIN	88 TO BE TAKEN UNTIL ALL USED	89 PHARMACIST WISHES TO SPEAK TO YOU	92 WARNING MAY CAUSE DEATH OR SERIOUS INJURY IF TAKEN BY ANOTHER PERSON.
95 PACK 1 USE THIS PACK FIRST	96 PACK 2 DO NOT USE THIS PACK UNTIL PACK 1 IS FINISHED	98 CYTOTOXIC	99 THIS MEDICINE MAY DECREASE THE EFFECTIVENESS OF ORAL CONTRACEPTIVES. CONSULT YOUR DOCTOR OR PHARMACIST.	100 A LOWER PRICED BRAND OF YOUR MEDICINE IS AVAILABLE PLEASE CONSULT YOUR DOCTOR OR PHARMACIST
102 PLEASE GIVE 48 HOURS NOTICE BEFORE NEXT SUPPLY.	108 PRESCRIPTION OWING FOR THIS ITEM	126 THIS PREPARATION IS AN ANTIBIOTIC. THE ENTIRE COURSE SHOULD BE COMPLETED.	127 SHAKE WELL AND KEEP IN REFRIGERATOR	

204 The Mixture SHAKE THE BOTTLE	205 CHILDREN'S PRESCRIPTION SHAKE THE BOTTLE	 <p>CYTOTOXIC</p> <p>HANDLE WITH CARE</p>		
200 POISON! Not to be taken	202 FOR EXTERNAL USE ONLY			
201 CAUTION! Not to be taken	203 SHAKE THE BOTTLE			

CAUTIONARY & ADVISORY LABEL HOLDERS



MC100



MC101